

Green Horse Wildlife References:

- Aney, W. C., & McClelland, B. R. (1990). Old-growth habitats and associated wildlife species in the northern Rocky Mountains (R1-90-42). Unpub. Report, USDA, Forest Service, Missoula MT.
- Arnett, E.B. & J.P. Hayes. 2009. Use of conifer snags as roosts by female bats in Western Oregon. *J. of Wildl. Mgmt.* 73(2):214-225.
- Badry, M. J. (2004). Fisher (*Martes pennanti*). Accounts and Measures for Managing Identified Wildlife—Accounts V. B.C. Min. of Water, Land and Air Protection. 19pp.
- Barclay, R.M.R. & A. Kurta. 2007. Ecology and behavior of bats roosting in tree cavities and under bark. in *Bats in Forests, Conservation and Management*. Eds. Lacki, M.J., J.P. Hayes & A. Kurta. pp. 17-59.
- Bartelt, P.E., C.R. Peterson, and R.W. Klaver. 2004. Sexual difference in the post-breeding movements and habitats selected by western toads (*Bufo boreas*) in southeastern Idaho. *Herpetologica* 60(4): 455-467.
- Bodie, W. L., Garton, E. O., Taylor, E. R., & McCoy, M. (1995). A sightability model for bighorn sheep in canyon habitats. *The Journal of Wildlife Management*, 832-840.
- Bonar, L. 2001. Pileated woodpecker habitat ecology in the Alberta foothills. Thesis, Univ. of Alberta, Edmonton, Alberta, Canada.
- Boyce, M. S., Mao, J. S., Merrill, E. H., Fortin, D., Turner, M. G., Fryxell, J., & Turchin, P. (2003). Scale and heterogeneity in habitat selection by elk in Yellowstone National Park. *Ecoscience*, 10(4), 421-431.
- Brewer, L.T., R. Bush, J.E. Canfield, & A.R. Dohmen. 2009. Northern goshawk northern region overview. USDAFS, Region 1, Missoula, MT. 54pp.
- Brown, S. R. (2015). Nez Perce-Clearwater National Forests VMap 2014. USDA Forest Service, Region 1. 9pp.
- Brown, S., & Barber, J. (2012). The Region 1 Existing Vegetation Mapping Program (VMAP) Flathead National Forest Overview (p. 5). Version 12. Rep. US Forest Service Region 1, 12-34.
- Brown, S., C. Hickey, B. Harrington, and R. Gill, eds. 2001. The U.S. Shorebird Conservation Plan, 2nd ed. Manomet Center for Conservation Sciences, Manomet, MA. 64pp.
- Browne, C. L. and C.A. Paszkowski. 2018. Microhabitat selection by western toads (*Anaxyrus boreas*). *Herp. Cons. & Biology* 13(2):317-330.
- Bull, E. L., Nielsen-Pincus, N., Wales, B. C., & Hayes, J. L. (2007). The influence of disturbance events on pileated woodpeckers in Northeastern Oregon. *Forest ecology and management*, 243(2-3), 320-329.
- Bull, E. L. 2006. Sexual differences in the ecology and habitat selection of western toads (*Bufo boreas*) in northeastern Oregon. Pacific Northwest Research Station La Grande, OR. *Herpetological Conservation and Biology* 1(1):27-38.

- Bull, E. L., Heater, T. W., & Shepherd, J. F. (2005). Habitat selection by the American marten in northeastern Oregon. *Northwest Science*. 79 (1): 36-42.
- Bull, E. L., and J. A. Jackson. 1995. Pileated woodpecker (*Dryocopus pileatus*). No. 148. A. Poole, and F. Gill, editors. The Academy of Natural Sciences, Philadelphia, Pennsylvania, and The American Ornithologists' Union, Washington, DC, USA.
- Bull, E. L., and R. S. Holthausen. 1993. Habitat use and management of pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management* 57: 335-345.
- Bull, E.L., S. Holthausen & M.G. Henjum. 1992. Roost trees used by pileated woodpeckers in Northeastern Oregon. *J. Wildl. Mgmt* V. 56(4):786-793.
- Bull, E. L., S.R. Peterson, and J.W. Thomas. 1986. Resource partitioning among woodpeckers in Northeastern Oregon. USDAFS, PNRS, PNW-444. Pp. 20
- Bush, Renate, D. Berglund, A. Leach, R. Lundberg, J.D. Zeiler. Overview of R1-FIA Summary Database, Region 1 Vegetation Classification, Inventory, and Analysis Report #06-02, 2006. http://fsweb.r1.fs.fed.us/forest/inv/fia_data/r1_fia_sdb.pdf
- Bush, R. and R. Lundberg. 2008. Wildlife habitat estimate update for the Region 1 Conservation Assessment. USDA Forest Service, Region 1, Missoula, MT. Numbered Report 08-04v1.0
- Busing, R. T., Halpern, C. B., & Spies, T. A. (1995). Ecology of Pacific yew (*Taxus brevifolia*) in western Oregon and Washington. *Conservation Biology*, 9(5), 1199-1207.
- Buskirk, S. W., & Ruggiero, L. F. (1994). American marten. In L. F. Ruggiero, K. B. Aubry, S. W. Buskirk, L. J. Lyon, & W. J. Zielinski (Eds.), The scientific basis for conserving forest carnivores: American marten, fisher, lynx, and wolverine (Gen. Tech. Rep. RM-254) (pp. 7-37). Fort Collins, CO: U.S. Department of Agriculture,, Forest Service, Rocky Mountain Forest and Range Experiment Station.
- Chapin, T., G., Harrison, D. J., & Katnik, D. D. (1998). Influence of landscape pattern on habitat use by American marten in an industrial forest. *Conservation Biology*, 12(6), 1327-1337.
- Christensen, A.G., L.J. Lyon & J.W. Unsworth. 1993. Elk management in the Northern Region: considerations in forest plan updates or revisions. USDAFS Intermountain Res. Sta. GTR INT-303.
- Cilimborg, A. 2006. Northern Region Landbird Monitoring Program: 2005 Flammulated owl surveys Final Report. Avian Science Center, U of MT, Missoula, MT. 18pp.
- Clapp, J. G., & Beck, J. L. (2016). Short-term impacts of fire-mediated habitat alterations on an isolated bighorn sheep population. *Fire Ecology*, 12(3), 80-98.
- Cook, J.G., L.L. Irwin, L.D. Bryant, R.A. Riggs & J.W. Thomas. 1998. Relations of forest cover and condition of elk: a test of thermal cover hypothesis in summer and winter. *Wildl. Monographs*, n 141:3-61.
- Crawford, R.C. 1983. Pacific Yew Community Ecology North-Central Idaho with implications to forest land management. Dissertation. U. of Idaho, Moscow, Idaho. 121 pp.

- Dibb, A. D., & Quinn, M. S. (2008). Response of bighorn sheep to restoration of Winter Range: revisited. In *Biennial Symposium of the Northern Wild Sheep and Goat Council* (Vol. 16, pp. 129-136). Cody, Wyoming, USA: Northern Wild Sheep and Goat Council.
- Dixon, R.D. and V.A. Saab. 2000. Black-backed woodpecker; in The Birds of North America, #509, American Ornith. Union, Wash. D.C.
- Dudley, J.G., Saab, V.A., & Hollenbeck, J.P. 2012. Foraging-habitat selection of Black-backed Woodpeckers in forest burns of southwestern Idaho. *The Condor*, 114(2), 348-357.
- Dudley, J. G., and V. A. Saab. 2007. Home range size of Blackbacked Woodpeckers in burned forests of southwestern Idaho. *Western North American Naturalist* 67:593–600
- Federal Register. 2011. Endangered and Threatened Wildlife and Plants; Reissuance of Final Rule To Identify the Northern Rocky Mountain Population of Gray Wolf as a Distinct Population Segment and To Revise the List of Endangered and Threatened Wildlife. V. 76, No. 87. 3pp.
- Federal Register. 2017. Endangered and Threatened Wildlife and Plants; 12-month Findings on Petitions to list 25 Species as Endangered or Threatened Species. (Fisher). Vol. 82, No. 192 on pp. 46634-46637.
- Federal Register. 2001. Executive Order 13186. Responsibilities of federal agencies to protect migratory birds. FR(v)66, No. 11. 1-17-2001. 3853-3856.
- Frair, J.L., E.H. Merrill, H.L. Beyer & J.M. Morales. 2008. Thresholds in landscape connectivity and mortality risks in response to growing road networks. *J. of Appl. Ecol.* 45:1504-1513.
- Giese, C.L., & F.J. Cuthbert, 2003. Influence of surrounding vegetation on woodpecker nest tree selection in oak forests of the Upper Midwest, USA. *For. Ecology & Mgmt*, 179(1), 523-534.
- Groves, C., Frederick, T., Frederick, G., Atkinson, E., Atkinson, M., Shepherd, J., & Servheen, G. (1997). Density, distribution, and habitat of Flammulated Owls in Idaho. *The Great Basin Naturalist*, 116-123.
- Guscio, C. G., B.R. Hossack, L.A. Eby, and P.S. Corn. 2007. Post-breeding habitat use by adult boreal toads (*Bufo boreas*) after wildfire in Glacier National Park, USA. *Herp. Cons. Biol.* 3 (1): 55-62.
- Gutiérrez, R. J. and David J. Delehanty. 1999. Mountain Quail (*Oreortyx pictus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/457>
- Hargis, C. D., Bissonette, J. A., & Turner, D. L. (1999). The influence of forest fragmentation and landscape pattern on American martens. *Journal of Applied Ecology*, 36(1), 157-172.
- Hayward, G. D.; Verner, J. 1994. Flammulated, boreal, and great gray owls in the United States: A technical conservation assessment. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 214 pp.
- Heekin, P.E. and K.P. Reese. 1995. Validation of a Mountain Quail Survey Technique. Tech. Rpt. FERC No. 1971; Hells Canyon Complex. U. of Idaho. 62 pp.

- Hoffman, N.J. 1997. Distribution of Picoides woodpeckers in relation to habitat disturbance within the Yellowstone area. Master Thesis, MSU, Bozeman, MT.
- Hoyt, J.S., & S.J. Hannon. 2002. Habitat associations of black-backed and three-toed woodpeckers in the boreal forest of Alberta. Can. J. of For. Res., 32(10), 1881-1888.
- Hutto, R. L., & Patterson, D. A. (2016). Positive effects of fire on birds may appear only under narrow combinations of fire severity and time-since-fire. *International Journal of Wildland Fire*, 25(10), 1074-1085.
- Hutto, R.L. 1995. Composition of bird communities following stand-replacement fires in northern Rock Mountain (USA) conifer forests. Cons. Biol. V.9(5):1041-1058.
- Idaho Department of Fish and Game. 2018. Moose. 76 pp. Boise, ID
- Idaho Department of Fish and Game. 2017. Wolf Statewide Report. 21 pp. Boise, ID
- Idaho Dept. of Fish and Game. 2016. Moose Report. 78 pp. Boise, ID.
- Idaho Department of Fish and Game (IDFG). 2014. Statewide furbearer progress report, surveys, and inventories. IDFG, Boise, ID
- Idaho Department of Fish and Game and Nez Perce Tribe. 2014. 2013 Idaho wolf monitoring progress report. Idaho Department of Fish and Game, 600 South Walnut, Boise, Idaho; Nez Perce Tribe Wolf Recovery Project, P.O. Box 365, Lapwai, Idaho. 74 pp.
- Idaho Department of Fish and Game (IDFG). 2010. Bighorn sheep management plan 2010. Idaho Department of Fish and Game, Boise, USA.
- Idaho Department of Fish and Game. 2005. Flammulated Owl. Webpage:
http://www.wildlife.uidaho.edu/idgap/idgap_report.asp).
- Idaho Department of Fish and Game. 2005. Fringed Myotis.
http://www.wildlife.uidaho.edu/idgap/idgap_report.asp).
- Idaho Department of Fish and Game. 2005. Mountain Quail.
http://www.wildlife.uidaho.edu/idgap/idgap_report.asp).
- Idaho Legislative Wolf Oversight Committee. 2002. Idaho wolf conservation and management plan. pp 32.
- Johnson, J. S., M. J. Lacki, and M. D. Baker. 2007. "Foraging ecology of long-legged myotis (*Myotis volans*) in north-central Idaho." Journal of Mammalogy 88, no. 5 (2007): 1261-1270.
- Keinath, D.A. 2004. Fringed Myotis (*Myotis thysanodes*): A technical conservation assessment. USDA Forest Service, Rocky Mountain Region. 64pp. Available:
<http://www.fs.fed.us/r2/projects/scp/assessments/fringedmyotis.pdf>
- Kennedy, P.L. (2003, January 2). Northern Goshawk (*Accipiter gentilis atricapillus*): a technical conservation assessment. [Online]. USDA Forest Service, Rocky Mountain Region. Available: <http://www.fs.fed.us/r2/projects/scp/assessments/northerngoshawk.pdf>

- Kirk, D. A., & Naylor, B. J. (1996). Habitat requirements of the Pileated Woodpecker (*Dryocopus pileatus*) with special reference to Ontario. Ontario Ministry of Natural Resources (No. 46, p. 49). SCST Technical Report.
- Koehler, G. M., & Hornocker, M. G. (1977). Fire effects on marten habitat in the Selway-Bitterroot Wilderness. *The Journal of Wildlife Management*, 500-505.
- Koehler, G. M., Moore, W. R., & Taylor, A. R. (1975). Preserving the pine marten: management guidelines for western forests. *Western Wildlands*.
- Kowalski, S. 2005. Frequency of Northern Goshawk Presence in the Northern Region 2005 Survey. USDA Forest Service, Region 1, Missoula, MT.
- Lacki, M. J., Baker, M. D., & Johnson, J. S. (2010). Geographic variation in roost-site selection of long-legged myotis in the Pacific Northwest. *The Journal of Wildlife Management*, 74(6), 1218-1228.
- Lacki, M. J., Johnson, J. S., Dodd, L. E., & Baker, M. D. (2007). Prey consumption of insectivorous bats in coniferous forests of north-central Idaho. *Northwest Science*, 81(3), 199-205.
- Lacki, M.J. & M.D. Baker. 2007. Day roosts of female fringed myotis (*Myotis thysanodes*) in xeric forests of the Pacific Northwest. *J. of Mammol.* 88(4):967-973.
- Linkhart, B.D. R.T. Reynolds and R.A. Ryder. 1998. Home range and habitat of breeding flammulated owls in Colorado. *T. Wilson Bull.* 110(3): 342-351.
- McKelvey, K.S., J.J. Claar, G.W. McDaniel and G. Hanvey. 1999. National lynx detection protocol. USDA Forest Service Forest Service, Rocky Mountain Research Station, Missoula, Montana, USA.
- Montgomery, R.A., G.J. Roloff & J.J. Millspaugh. 2012. Variation in elk response to roads by season, sex and road type. *J. of Wildl. Mgmt.* 77(2):313-325.
- Moriarty, K. M., Zielinski, W. J., & Forsman, E. D. (2011). Decline in American marten occupancy rates at Sagehen Experimental Forest, California. *The Journal of Wildlife Management*, 75(8), 1774- 1787.
- Moser, B. W., & Garton, E. O. (2019). Northern goshawk space use and resource selection. *The Journal of Wildlife Management*, 83(3), 705-713.
- Moser, B. W., & Garton, E. O. (2009). Short-term effects of timber harvest and weather on Northern Goshawk reproduction in northern Idaho. *Journal of Raptor Research*, 43(1), 1-11.
- Moser, B.W. 2007. Space use and ecology of goshawks in Northern Idaho. Dissertation in Natural Resources, U of I, Moscow, Idaho.
- Murphy, E., & W. Lehnhausen. 1998. Density and foraging ecology of woodpeckers following a stand-replacement fire. *J. of Wildl. Mgmt.*, 62(4), 1359-1372.
- Nadeau, M. S., DeCesare, N. J., Brimeyer, D. G., Bergman, E. J., Harris, R. B., Hersey, K. R., ... & Thomas, T. P. (2017). Status and trends of moose populations and hunting opportunity in

- the western United States. *Alces: A Journal Devoted to the Biology and Management of Moose*, 53, 99-112.
- Nappi, A., Drapeau, P., Saint-Germain, M., & Angers, V. A. (2010). Effect of fire severity on long-term occupancy of burned boreal conifer forests by saproxylic insects and wood-foraging birds. *International Journal of Wildland Fire*, 19(4), 500-511.
- NatureServe. 2019. Online database that tracks species status primarily in North America.
- Nez Perce-Clearwater National Forest. 2018. Monitoring report 2013-2017. Pp.16.
- Nez Perce-Clearwater National Forests Plan Revision. Draft Environmental Impact Statement. Chapter 3, 3.2.3. Wildlife pp. 83-88.
- Olson, L. E., Sauder, J. D., Albrecht, N. M., Vinkey, R. S., Cushman, S. A., & Schwartz, M. K. (2014). Modeling the effects of dispersal and patch size on predicted fisher (Pekania [Martes pennanti]) distribution in the US Rocky Mountains. *Biological Conservation*, 169, 89-98.
- Pierce, D. J., & Peek, J. M. (1984). Moose habitat use and selection patterns in north-central Idaho. *The Journal of wildlife management*, 1335-1343.
- Poole, K. G., & Stuart-Smith, K. (2005). Finescale winter habitat selection by moose in interior montane forests. *Alces*, 41(1), 1-8.
- Potvin, F., Bélanger, L., & Lowell, K. (2000). Marten habitat selection in a clearcut boreal landscape. *Conservation Biology*, 14(3), 844-857.
- Powers, L.R., A. Dale, P.A. Gaede, C. Rodes, L. Nelson, J.J. Dean, J.D. May. 1996/ Nesting and food habits of the flammulated owl (Otus flammeolus) in southcentral Idaho. *J.Raptor Res.* 30(1): 15-20.
- Raley, C.M., E.C. Lofroth, R.L. Truex, J.S. Yaeger, & J.M. Higley. 2012. Habitat ecology of fishers in Western North America. In *Biology and Conservation of martens, sables and fishers*. Editors: Aubry, Zielinski, Raphael, Proulx,& Buskirk. Chapter 10: 231-254.
- Ranglack, D.H, K.M. Proffitt, J. Gude, J. Canfield, J. Rotella, & R.A. Garrott. 2017. Security areas for elk during archery and rifle hunting seasons. *J. Wldlf Mgmt*. v.81(5):778-791.
- Reynolds, R.T. and B.D Linkhart. 1992. Flammulated owls in ponderosa pine: evidence of preference for old growth. In *MR Kaufman, WH Moir, and RL Bassett, technical coordinators. Proceedings of the workshop on old-growth in the Southwest and Rocky Mountain Region. Portal, Arizona, USA* (pp. 166-169).
- Rich, T.D., 2007. Guide to the Partners in Flight Population Estimates Database. Version: North American Landbird Conservation Plan 2004. Partners in Flight Technical Series No 5. <http://www.partnersinflight.org/>
- Ritchie, M. W., Knapp, E. E., & Skinner, C. N. (2013). Snag longevity and surface fuel accumulation following post-fire logging in a ponderosa pine dominated forest. *Forest Ecology and Management*, 287, 113-122.
- Romin, S.J. and W.R. Bosworth. 2010. Northern Region Bats Species Occurrences in Idaho. IDFG & USFS. Missoula, MT. Agreement No. AG-03R6-P-08-0251. 69pp.

- Rowland, M.M., M.J. Wisdom, B.K. Johnson & M.A. Penninger. 2005. Effect of roads on elk: implications for management in forested ecosystems. Pages 42-52 in Wisdom, M. J., technical editor, The Starkey Project: a synthesis of long-term studies of elk and mule deer.
- Rowland, M.M., M.J. Wisdom, B.K. Johnson & J.G. Kie. 2000. Elk distribution and modeling in relation to roads. *J. of Wildl. Mgmt.* v.64(3):672-684.
- Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon and W.J. Zielinski (tech. eds.). 1994. The Scientific basis for conserving forest carnivores: American marten, fisher, lynx and wolverine in the western United States. General Technical Report RM-GTR-254. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 184 pp.
- Russell, R. E., Saab, V. A., Rotella, J. J., & Dudley, J. G. (2009). Detection probabilities of woodpecker nests in mixed conifer forests in Oregon. *The Wilson Journal of Ornithology*, 121(1), 82-88.
- Sauder, J.D. and J.L. Rachlow. 2015. Forest heterogeneity influences habitat selection by fishers (Pekania pennant) within home ranges. *For. Ecol. & Mgmt.* 347: 49-56.
- Sauder, J.D. 2014. Landscape ecology of fishers (Pekania pennant) in North-Central Idaho. Dissertation in Natural Resources. U of Idaho, Moscow, Idaho.
- Sauder, J.D. and J.L. Rachlow. 2014. Both forest composition and configuration influence landscape-scale habitat selection by fishers (Pekania pennanti) in mixed coniferous forests of the Northern Rocky Mountains. *Forest Ecology and Mgmt.* 314: 75-84.
- Schremp, T.V. 2017. Diet selection, forage, quality, and forage availability: could forage limit moose populations in Northern Idaho? Thesis. U of I, Moscow, ID. 71pp.
- Schultz, C. 2010. Effective wildlife conservation planning. *BioScience.*, V. 60, N.7, 545-551.
- Schwartz, M. K., DeCesare, N. J., Jimenez, B. S., Copeland, J. P., & Melquist, W. E. (2013). Stand-and landscape-scale selection of large trees by fishers in the Rocky Mountains of Montana and Idaho. *Forest Ecology and Management*, 305, 103-111.
- Servheen, G. 1997. Interagency guidelines for evaluating and managing elk habitats and populations in Central Idaho. 67pp.
- Shirk, A. J., Raphael, M. G., & Cushman, S. A. (2014). Spatiotemporal variation in resource selection: insights from the American marten (*Martes americana*). *Ecological Applications*, 24(6), 1434-1444.
- Smith, T. S., Hardin, P. J., & Flinders, J. T. (1999). Response of bighorn sheep to clear-cut logging and prescribed burning. *Wildlife Society Bulletin*, 840-845.
- Snyder, J. E., & Bissonette, J. A. (1987). Marten use of clear-cuttings and residual forest stands in western Newfoundland. *Canadian Journal of Zoology*, 65(1), 169-174.
- Squires, J. R., and P. L. Kennedy. 2006. Northern goshawk ecology: an assessment of current knowledge and information needs for conservation management. *Studies in Avian Biology* 31: 8-62.

- Squires, J. R., & Reynolds, R. T. (1997). Northern goshawk (*Accipiter gentilis*). In A. Poole & F. Gill (Eds.), *The Birds of North America*, No. 298 (pp. 43). Washington, DC: The Academy of Natural Sciences, The American Ornithologists' Union.
- Stickney, P.F. 1980. Data base for post-fire succession, first 6 to 9 years, in Montana larch-fir forests. USDA Forest Service. Gen. Tech Rpt. 148 pp.
- Thomas, J.W., H. Black, R.J. Scherzinger & R.J. Pedersen. 1979. Deer and Elk, in Wildlife habitats in managed forests, the Blue Mountains of Oregon and Washington. 104-127.
- Tremblay, J., Ibarzabal, J., Dussault, C., & Savard, J. P. (2009). Habitat requirements of breeding Black-backed Woodpeckers (*Picoides arcticus*) in managed, unburned boreal forest. *Avian Conservation and Ecology*, 4(1).
- Unsworth, J.W., L. Kuck, E.O. Garton, and B.R. Butterfield. 1998. Elk habitat selection on the Clearwater National Forest, Idaho. *Journal of Wildlife Management*. 62(4): 1255-1263.
- Unsworth, J.W., L. Kuck, M.D. Scott, & E.O. Garton. 1993. Elk mortality in the Clearwater drainage of Northcentral Idaho. *J. Wildl. Mgmt* 57(3):495-502.
- USDA Forest Service. 2008. Memorandum of understanding between the U.S. Department of Agriculture Forest Service and the U.S. Fish and Wildlife Service to promote the conservation of migratory birds. FS Agreement #08-MU-1113-2400-264. 13pp.
- USDI Fish and Wildlife Service. 1998. Northern Goshawk Status Review. US Fish and Wildlife Service, Portland, Oregon, USA
- USDI Fish and Wildlife Service. 1987. Northern Rocky Mountain Wolf Recovery Plan. Denver, CO. 119pp.
- USDI Fish and Wildlife Service. 2013. 50 CFR Part 17: Endangered and Threatened Wildlife and Plants; Revised Designation of Critical Habitat for the Contiguous U.S. Distinct Population Segment of the Canada Lynx and Revised Distinct Population Segment Boundary; Proposed Rule. FR, v.78, n 187.
- US Presidential Executive Order 13186. 2001. Federal Register (V-66, No. 11, pages 3853-3856.
- Vonhof and Barclay. 1997. Use of tree stumps as roosts by the western long-eared bat. *J. Wildl. Mgmt.* 61(3):674-684.
- Wagner, G. D., & Peek, J. M. (2006). Bighorn sheep diet selection and forage quality in central Idaho. *Northwest Science*, 80(4), 246.
- Wasserman, T. N., Cushman, S. A., Shirk, A. S., Landguth, E. L., & Littell, J. S. (2012). Simulating the effects of climate change on population connectivity of American marten (*Martes americana*) in the northern Rocky Mountains, USA. *Landscape Ecology*, 27(2), 211-225.
- Wasserman, T. N., Cushman, S. A., Schwartz, M. K., & Wallin, D. O. (2010). Spatial scaling and multi-model inference in landscape genetics: *Martes americana* in northern Idaho. *Landscape ecology*, 25(10), 1601-1612.

- Webb, S. M. (2009). Marten fur harvests and landscape change in west-central Alberta. *The Journal of Wildlife Management*, 73(6), 894-903.
- Weir, R.D. & A.S. Harestad. 2003. Scale dependent habitat selectivity by fishers in South-Central British Columbia. *J. of Wildl. Mgmt.* 67(1):73-82.
- Weir, R. D., & Corbould, F. B. (2010). Factors affecting landscape occupancy by fishers in north-central British Columbia. *Journal of Wildlife Management*, 74(3), 405-410.
- Wisdom, M.J, R.S. Holthausen, B.C. Wales, C.D. Hargis, V.A. Saab, D.C. Lee, W.J. Hahn, T.D. Rich, M.M. Rowland, W.J. Murphy, & M.R. Eames. 2000. Source habitats for terrestrial vertebrates, of focus in the interior Columbia Basin: broad-scale trends and management implications. Volume 1.